INFORMATION DISCLOSURE STATEMENT BY APPLICANT

(Not for submission under 37 CFR 1.99)

		Approved for use through 07/31/2012.		
	Application Number	10/574,477		
	Confirmation Number	2361		
	Filing Date	January 9, 2007		
	First Named Inventor	Masao SUDOH		
	Art Unit	1621		
ı	Examiner Name	Sudhakar KATAKAM		
	Attorney Docket Number	Q94121		

U.S. PATENTS							
Examiner Initials*	Cite No	Patent Number	Kind Code ¹	Issue Date	Name of Patentee or Applicant of cited Document	Pages, Columns, Lines, Where Relevant Passages or Relevant Figures Appear	

	U.S. PATENT APPLICATION PUBLICATIONS						
Examiner Initials*	Cite No	Publication Number	Kind Code ¹	Publication Date	Name of Patentee or Applicant of cited Document	Pages, Columns, Lines, Where Relevant Passages or Relevant Figures Appear	

	FOREIGN PATENT DOCUMENTS							
Examiner Initials*	Cite No	Foreign Document Number ³	Country Code ²	Kind Code ⁴	Publication Date	Name of Patentee or Applicant of cited Document	Pages, Columns, Lines, Where Relevant Passages or Relevant Figures Appear	T⁵

NON-PATENT LITERATURE DOCUMENTS						
Examiner Initials* Cite No Cite No Cite No United name of the author (in CAPITAL LETTERS), title of the article (when appropriate), title of the item (book, magazine, journal, serial, symposium, calatiog, etc.), date, page(s), volume-issue number(s), published.						
	1.	Japanese Office Action issued on October 1, 2010 in the corresponding Japanese Patent Application No. 2005-514507	Υ			

	EXAMINER SIGNATURE					
Examiner Signature		Date Considered				

*EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609. Draw line through a citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

1 See Kin Codes of USPTO Patient Documents at www.USPTO.GOV or MPEP.901.04. 2 Either office in this issued the document, by the two-letter code (WIPO Standard at 785). 3 For Japones period focuments, the indication of the year of the region of the Empirer or patient percent be sensited by the percent in the patient of t